arXiv:1201.0141v1 [math.PR] 30 Dec 2011

HIGHER-ORDER LAPLACE EQUATIONS AND HYPER-CAUCHY DISTRIBUTIONS

ENZO ORSINGHER AND MIRKO D'OVIDIO

Abstract. In this paper we introduce new distributions which are solutions of higher-order Laplace equations. It is proved that their densities can be obtained by folding and symmetrizing Cauchy distributions. Another class of probability laws related to higher-order Laplace equations is obtained by composing pseudo-processes with positively-skewed Cauchy distributions which produce asymmetric Cauchy densities in the odd-order case. A special attention is devoted to the third-order Laplace equation where the connection between the Cauchy distribution and the Airy functions is obtained and analyzed.

1. Introduction

The Cauchy density

$$p(x,t) = \frac{1}{\pi} \frac{t}{(x^2+t^2)} \tag{1.1}$$

solves the Laplace equation (see Nane [8])

$$\frac{\partial^2 u}{\partial t^2} + \frac{\partial^2 u}{\partial x^2} = 0, \quad x \in \mathbb{R},\ t > 0. \tag{1.2}$$

The n-dimensional counterpart of (1.1)

$$p(\mathbf{x},t) = \frac{\Gamma\left(\frac{n}{2}\right)}{\pi^{\frac{n}{2}}} \frac{t}{(t^2+|\mathbf{x}|^2)^{\frac{n}{2}}}, \quad \mathbf{x} \in \mathbb{R}^{n-1},\ t > 0 \tag{1.3}$$

with characteristic function

$$\int_{\mathbb{R}^{n-1}} e^{i\langle \boldsymbol{\alpha}, \mathbf{x}\rangle} p(\mathbf{x},t) d\mathbf{x} = \exp\left(-t|\boldsymbol{\alpha}|\right) \tag{1.4}$$

solves the n-dimensional Laplace equation

$$\frac{\partial^2 p}{\partial t^2} + \sum_{j=1}^{n-1} \frac{\partial^2 p}{\partial x_j^2} = 0. \tag{1.5}$$

The inspiring idea of this paper is to investigate the class of distributions which satisfy the higher-order Laplace equations of the form

$$\frac{\partial^n u}{\partial t^n} + \frac{\partial^n u}{\partial x^n} = 0, \quad x \in \mathbb{R},\ t > 0 \tag{1.6}$$

Date: January 4, 2012.

2010 *Mathematics Subject Classification.* 60G52,35C05.

Key words and phrases. Pseudo-processes, stable processes, Cauchy processes, higher-order Laplace equations, Airy functions, modified Bessel functions.

In a previous paper of ours we have shown that the law

$$p_4(x,t) = \frac{t}{\pi\sqrt{2}}\frac{x^2+t^2}{x^4+t^4} \tag{1.7}$$

solves the fourth-order Laplace equation

$$\frac{\partial^4 u}{\partial t^4} + \frac{\partial^4 u}{\partial x^4} = 0, \quad x \in \mathbb{R},\ t > 0. \tag{1.8}$$

In Section 2 we analyze distributions related to equations of the form

$$\frac{\partial^{2^n} u}{\partial t^{2^n}} + \frac{\partial^{2^n} u}{\partial x^{2^n}} = 0 \tag{1.9}$$

which can be expressed in many alternative forms. The decoupling of the 2^n-th order differential operator in (1.9)

$$\frac{\partial^{2^n}}{\partial t^{2^n}} + \frac{\partial^{2^n}}{\partial x^{2^n}} = \prod_{\substack{k=-(2^{n-1}-1)\\ k\ odd}}^{2^{n-1}-1} \left(\frac{\partial^2}{\partial t^2} + e^{i\frac{\pi k}{2^{n-1}}} \frac{\partial^2}{\partial x^2} \right)$$

suggests to represent distributions related to (1.9) as

$$p_{2^n}(x,t) = \frac{1}{\pi 2^{n-1}} \sum_{\substack{k=-(2^{n-1}-1)\\ k\ odd}}^{2^{n-1}-1} \frac{t\, e^{i\frac{\pi k}{2^n}}}{x^2 + (te^{i\frac{\pi k}{2^n}})^2}, \quad n \geq 2. \tag{1.10}$$

that is the superposition of Cauchy densities at imaginary times. Alternatively, we give a real-valued expression for (1.10) as

$$p_{2^n}(x,t) = \frac{t(x^2+t^2)}{2^{n-2}\pi(x^{2^n}+t^{2^n})} \sum_{\substack{k=1\\ k \text{ odd}}}^{2^{n-1}-1} \prod_{\substack{j=1,\, j \text{ odd}\\ j\neq k}}^{2^{n-1}-1} \left(x^4 + t^4 + 2x^2t^2 \cos\frac{j\pi}{2^{n-1}} \right). \tag{1.11}$$

The density (1.11) can also be represented as

$$p_{2^n}(x,t) = \frac{t(x^2+t^2)}{2^{n-2}\pi} \sum_{\substack{k=1\\ k \text{ odd}}}^{2^{n-1}-1} \frac{\cos\frac{k\pi}{2^n}}{x^4+t^4+2x^2t^2\cos\frac{k\pi}{2^{n-1}}}, \quad n \geq 2. \tag{1.12}$$

Each component of the distribution (1.12) is produced by folding and symmetrizing the density of the r.v.

$$V(t) = C\left(t\cos\frac{k\pi}{2^n}\right) - t\sin\frac{k\pi}{2^n}, \quad t>0,\ 1 \leq k \leq 2^{n-1}-1,\ k \text{ odd}$$

where $C(t)$, $t > 0$ is the Cauchy symmetric process. The distributions (1.12) differ from the Cauchy laws since they have a bimodal structure for all $n \geq 2$ as figures below show. For $n = 2$, the distribution (1.11) reduces to (1.7) if we assume that the inner product appearing in formula (1.11) is equal to one. Of course, the density (1.12) coincides with (1.7) for $n = 2$. For $n = 3$ we get from (1.11) and (1.12) that

$$\begin{aligned} p_{2^3}(x,t) =& \frac{t(x^2+t^2)}{\sqrt{2}\pi(x^8+t^8)} \left[(x^4+t^4-\sqrt{2}x^2t^2)\cos\frac{\pi}{8} + (x^4+t^4+\sqrt{2}x^2t^2)\sin\frac{\pi}{8} \right] \\ =& \frac{t(x^2+t^2)}{2\pi} \left[\frac{\sin\frac{\pi}{8}}{x^4+t^4-\sqrt{2}x^2t^2} + \frac{\cos\frac{\pi}{8}}{x^4+t^4+\sqrt{2}x^2t^2} \right]. \end{aligned} \tag{1.13}$$

In Orsingher and D'Ovidio [11] we have shown that the density (1.7) is the probability distribution of

$$Q(t) = F(T_t), \quad t > 0$$

where F is the Fresnel pseudo-process described in [11] and T_t, $t > 0$ is the first passage time of a Brownian motion independent from F. We note that

$$\mathcal{Q}(t) = F(|B(t)|), \quad t > 0$$

has density coinciding with the fundamental solution of the fourth-order heat equation

$$\frac{\partial u}{\partial t} = -\frac{\partial^4 u}{\partial x^4}.$$

We prove also that, for $k \in \mathbb{N}$, there are non-centered Cauchy distributions which solve the equations

$$\frac{\partial^{2k+1} u}{\partial t^{2k+1}} + \frac{\partial^{2k+1} u}{\partial x^{2k+1}} = 0. \tag{1.14}$$

If $X_{2k+1}(t)$, $t > 0$ is the pseudo-process whose density measure

$$\mu_{2k+1}(dx, t) = \mu\{X_{2k+1}(t) \in dx\}$$

solves the heat-type equations

$$\frac{\partial u}{\partial t} = -\frac{\partial^{2k+1} u}{\partial x^{2k+1}}, \quad k \in \mathbb{N} \tag{1.15}$$

and $S_{\frac{1}{2k+1}}(t)$, $t > 0$ is a positively skewed stable process of order $\frac{1}{2k+1}$ we have that

$$Pr\{X_{2k+1}(S_{\frac{1}{2k+1}}(t)) \in dx\}/dx = \frac{t \cos \frac{\pi}{2(2k+1)}}{\pi \left[\left(x + (-1)^{k+1} t \sin \frac{\pi}{2(2k+1)} \right)^2 + t^2 \cos^2 \frac{\pi}{2(2k+1)} \right]}. \tag{1.16}$$

We show below that the densities (1.16) solve also the following second-order p.d.e.

$$\frac{\partial^2 u}{\partial t^2} + \frac{\partial^2 u}{\partial x^2} = 2 \sin \frac{\pi}{2(2k+1)} \frac{\partial^2 u}{\partial t \, \partial x}.$$

We have investigated in detail the case of third-order Laplace equation

$$\frac{\partial^3 u}{\partial t^3} + \frac{\partial^3 u}{\partial x^3} = 0 \tag{1.17}$$

and have shown that

$$\begin{aligned} Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\} =& dx \int_0^\infty \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{x}{\sqrt[3]{3s}}\right) \frac{t}{s} \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{t}{\sqrt[3]{3s}}\right) ds \qquad (1.18) \\ =& dx \frac{\sqrt{3}}{2} t \frac{x-t}{x^3 - t^3} = dx \frac{\sqrt{3}}{2} \frac{t}{x^2 + xt + t^2} \\ =& dx \frac{t \cos \frac{\pi}{6}}{(x + t \sin \frac{\pi}{6})^2 + t^2 \cos^2 \frac{\pi}{6}}. \end{aligned}$$

The pictures of the Cauchy distributions (1.16) show that the location parameter $t \sin \frac{\pi}{2(2k+1)}$ tends to zero as $k \to \infty$ while the scale parameter tends to one, $t \cos \frac{\pi}{2(2k+1)} \to t$. This means that the asymmetry of the Cauchy densities decreases as k increases and is maximal for $k = 1$. The decrease of parameters of (1.16) (with k increasing) is due to the growing symmetrization of the fundamental solutions of equations (1.15).

By suitably combining the distribution (1.16) for $k = 1$, we arrive at the density

$$p_6(x,t) = \frac{\sqrt{3}}{2^2\pi} t \frac{(x^2+t^2)\cos\frac{\pi}{6} + xt}{(x^2+t^2+xt\cos\frac{\pi}{6})^2 + 2x^2t^2\cos\frac{\pi}{3}} \tag{1.19}$$

which solves the equation

$$\frac{\partial^6 u}{\partial t^6} + \frac{\partial^6 u}{\partial x^6} = 0. \tag{1.20}$$

The probability density (1.19) displays the unimodal structure of the Cauchy distribution.

2. Hyper Cauchy distributions

In this section we analyze the distribution related to Laplace-type equations of the form

$$\left(\frac{\partial^{2^n}}{\partial t^{2^n}} + \frac{\partial^{2^n}}{\partial x^{2^n}} \right) u = 0, \quad n > 1. \tag{2.1}$$

For $n \geq 2$ we obtain a new class of distributions having the form

$$p_{2^n}(x,t) = \frac{t(x^2+t^2)}{2^{n-2}\pi(x^{2^n}+t^{2^n})} g(x,t), \quad x \in \mathbb{R},\ t > 0 \tag{2.2}$$

where $g(x,t)$ is a polynomial of order $2^n - 2^2$. For $n = 2$, formula (2.2) yields the distribution

$$p_4(x,t) = \frac{t(x^2+t^2)}{\sqrt{2}\pi(x^4+t^4)}, \quad x \in \mathbb{R},\ t > 0 \tag{2.3}$$

emerging in the analysis of Fresnel pseudo-processes (see Orsingher and D'Ovidio [11]).

The main result of this section is given in the next theorem.

Theorem 2.1. *The hyper Cauchy density*

$$p_{2^n}(x,t) = \frac{1}{\pi 2^{n-1}} \sum_{\substack{k=-(2^{n-1}-1) \\ k \text{ odd}}}^{2^{n-1}-1} \frac{t e^{i\frac{\pi k}{2^n}}}{x^2 + (t e^{i\frac{\pi k}{2^n}})^2} \tag{2.4}$$

solves the equation

$$\left(\frac{\partial^{2^n}}{\partial t^{2^n}} + \frac{\partial^{2^n}}{\partial x^{2^n}} \right) u = 0, \quad x \in \mathbb{R},\ t > 0, \quad n > 1. \tag{2.5}$$

A real-valued expression of (2.4) *reads*

$$p_{2^n}(x,t) = \frac{t(x^2+t^2)}{\pi 2^{n-2}(x^{2^n}+t^{2^n})} \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-1}-1} \cos\frac{k\pi}{2^n} \prod_{\substack{k \neq j=1 \\ j \text{ odd}}}^{2^{n-1}-1} \left(x^4 + t^4 + 2x^2t^2\cos\frac{j\pi}{2^{n-1}} \right) \tag{2.6}$$

or equivalently

$$p_{2^n}(x,t) = \frac{t(x^2+t^2)}{2^{n-2}\pi} \sum_{\substack{k=1 \\ k \text{ odd}}}^{2^{n-1}-1} \frac{\cos\frac{k\pi}{2^n}}{x^4+t^4+2x^2t^2\cos\frac{k\pi}{2^{n-1}}}, \quad \textit{for } n > 1. \tag{2.7}$$

Proof. In order to check that (2.4) satisfies equation (2.5) we resort to Fourier transforms

$$U(\beta,t)=\int_{-\infty}^{+\infty}e^{i\beta x}u(x,t)dx.$$

Equation (2.5) becomes

$$\frac{\partial^{2^n}U}{\partial t^{2^n}}+(-i\beta)^{2^n}U=\frac{\partial^{2^n}U}{\partial t^{2^n}}+\beta^{2^n}U=0. \tag{2.8}$$

The solutions of the algebraic equation associated to (2.8) have the form

$$r_j=|\beta|\,e^{i\pi\frac{2j+1}{2^n}},\quad 0\le j\le 2^n-1. \tag{2.9}$$

In order to construct bounded solutions to (2.8) we restrict ourselves to

$$U(\beta,t)=\frac{1}{2^{n-1}}\sum_{\substack{k=-(2^{n-1}-1)\\ k\ odd}}^{2^{n-1}-1}e^{-t|\beta|e^{i\frac{k\pi}{2^n}}} \tag{2.10}$$

where the normalizing constant in (2.10) is chosen equal to $1/2^{n-1}$ so that $U(\beta,0)=1$. The inverse of (2.10) is (2.4). We check directly that each term of (2.4) has Fourier transform solving equation (2.8). For all odd values of k, we have that

$$\begin{aligned}&\int_{-\infty}^{+\infty}e^{i\beta x}\left(\frac{\partial^{2^n}}{\partial t^{2^n}}+\frac{\partial^{2^n}}{\partial x^{2^n}}\right)\left(\frac{te^{i\frac{k\pi}{2^n}}}{x^2+(te^{i\frac{k\pi}{2^n}})^2}\right)dx\\ =&\frac{\partial^{2^n}}{\partial t^{2^n}}e^{-t|\beta|e^{i\frac{k\pi}{2^n}}}+(-i\beta)^{2^n}e^{-t|\beta|e^{i\frac{k\pi}{2^n}}}\\ =&\left(\beta^{2^n}e^{ik\pi}+i^{2^n}\beta^{2^n}\right)e^{-t|\beta|e^{i\frac{k\pi}{2^n}}}\\ =&\left((-1)^k\beta^{2^n}+\beta^{2^n}\right)e^{-t|\beta|e^{i\frac{k\pi}{2^n}}}=0\end{aligned}$$

because k is odd. In order to obtain (2.6) we observe that, in view of (2.4) we can write

$$p_{2^n}(x,t)=\frac{1}{\pi}\sum_{\substack{k=-(2^{n-1}-1)\\ k\text{ odd}}}^{2^{n-1}-1}\frac{c_k\,t^{|2k-1|}x^{2^n-|2k-1|-1}}{\prod_{\substack{k=-(2^{n-1}-1)\\ k\text{ odd}}}^{2^{n-1}-1}(x^2+(te^{i\frac{k\pi}{2^n}})^2)}$$

where

$$\prod_{\substack{k=-(2^{n-1}-1)\\ k\text{ odd}}}^{2^{n-1}-1}(x^2+(te^{i\frac{k\pi}{2^n}})^2)=x^{2^n}+t^{2^n} \tag{2.11}$$

and c_k are constants evaluated below. Result (2.11) can be obtained directly by solving the equation $x^{2^n}+t^{2^n}=0$ or by successively regrouping the terms of the right-hand side of (2.11). We have at first that

$$\prod_{\substack{k=-(2^{n-1}-1)\\ k\text{ odd}}}^{2^{n-1}-1}(x^2+(te^{i\frac{k\pi}{2^n}})^2)=\prod_{k=1,\,k\text{ odd}}^{2^{n-1}-1}\left(x^4+t^4+2x^2t^2\cos\frac{k\pi}{2^{n-1}}\right)$$

$$
\begin{aligned}
&=\prod_{k=1,\, k \text{ odd}}^{2^{n-2}-1}\left(x^8+t^8+2x^4t^4\cos\frac{k\pi}{2^{n-2}}\right)\\
&=\cdots\\
&=\left(x^{2^n}+t^{2^n}+2x^2t^2\cos\frac{\pi}{2}\right)\\
&=x^{2^n}+t^{2^n}.
\end{aligned}
$$

In view of (2.11) we can rewrite (2.4) as

$$
p_{2^n}(x,t)=\frac{t}{\pi 2^{n-1}(x^{2^n}+t^{2^n})}\sum_{\substack{k=-(2^{n-1}-1)\\ k \text{ odd}}}^{2^{n-1}-1}\ \prod_{\substack{j=-(2^{n-1}-1)\\ j \text{ odd},\, j\neq k}}^{2^{n-1}-1}(x^2+(te^{i\frac{2\pi j}{2^n}}))e^{i\frac{\pi k}{2^n}}
$$

where

$$
\begin{aligned}
&\sum_{\substack{k=-(2^{n-1}-1)\\ k \text{ odd}}}^{2^{n-1}-1}\ \prod_{\substack{j=-(2^{n-1}-1)\\ j \text{ odd},\, j\neq k}}^{2^{n-1}-1}(x^2+(te^{i\frac{2\pi j}{2^n}}))e^{i\frac{\pi k}{2^n}}\\
=&\sum_{\substack{k=-(2^{n-1}-1)\\ k \text{ odd}}}^{2^{n-1}-1}\ \prod_{\substack{j=1\\ j \text{ odd},\, j\neq k}}^{2^{n-1}-1}(x^4+t^4+2x^2t^2\cos\frac{\pi j}{2^{n-1}})(x^2+(te^{-i\frac{2\pi k}{2^n}}))e^{i\frac{\pi k}{2^n}}\\
=&\sum_{\substack{k=1\\ k \text{ odd}}}^{2^{n-1}-1}\ \prod_{\substack{j=1\\ j \text{ odd},\, j\neq k}}^{2^{n-1}-1}(x^4+t^4+2x^2t^2\cos\frac{\pi j}{2^{n-1}})(x^2e^{i\frac{k\pi}{2^n}}+t^2e^{-i\frac{k\pi}{2^n}}+x^2e^{-i\frac{k\pi}{2^n}}+t^2e^{i\frac{k\pi}{2^n}})\\
=&2(x^2+t^2)\sum_{\substack{k=1\\ k \text{ odd}}}^{2^{n-1}-1}\cos\frac{k\pi}{2^n}\prod_{\substack{j=1\\ j \text{ odd},\, j\neq k}}^{2^{n-1}-1}(x^4+t^4+2x^2t^2\cos\frac{\pi j}{2^{n-1}})
\end{aligned}
$$

and thus

$$
p_{2^n}(x,t)=\frac{t(x^2+t^2)}{\pi 2^{n-2}(x^{2^n}+t^{2^n})}\sum_{\substack{k=1\\ k \text{ odd}}}^{2^{n-1}-1}\cos\frac{k\pi}{2^n}\prod_{\substack{j=1\\ j \text{ odd},\, j\neq k}}^{2^{n-1}-1}(x^4+t^4+2x^2t^2\cos\frac{\pi j}{2^{n-1}}).
$$

Furthermore, from the fact that

$$
x^{2^n}+t^{2^n}=\prod_{k=1,\, k \text{ odd}}^{2^{n-2}-1}\left(x^8+t^8+2x^4t^4\cos\frac{k\pi}{2^{n-2}}\right)
$$

we obtain that

$$
p_{2^n}(x,t)=\frac{t(x^2+t^2)}{2^{n-2}\pi}\sum_{\substack{k=1\\ k \text{ odd}}}^{2^{n-1}-1}\frac{\cos\frac{k\pi}{2^n}}{x^4+t^4+2x^2t^2\cos\frac{k\pi}{2^{n-1}}}.
$$

$\square$

Remark 2.2. In order to prove that the density (2.6) integrates to unity we present the following calculation

$$\begin{aligned}
\int_{-\infty}^{+\infty} \frac{x^2+t^2}{x^4+t^4+2x^2t^2\cos\frac{k\pi}{2^{n-1}}}dx =& 2\int_0^{+\infty} \frac{x^2+t^2}{x^4+t^4+2x^2t^2\cos\frac{k\pi}{2^{n-1}}}dx \\
=& \frac{2}{t}\int_0^{+\infty} \frac{y^2+1}{y^4+1+2y\cos\frac{k\pi}{2^{n-1}}}dy \\
=& \frac{2}{t}\int_0^{\frac{\pi}{2}} \frac{1}{\tan^4\theta+1+2\tan^2\theta\cos\frac{k\pi}{2^{n-1}}}\frac{d\theta}{\cos^4\theta} \\
=& \frac{2}{t}\int_0^{\frac{\pi}{2}} \frac{d\theta}{\sin^4\theta+\cos^4\theta+2\sin^2\theta\cos^2\theta\cos\frac{k\pi}{2^{n-1}}} \\
=& \frac{2}{t}\int_0^{\frac{\pi}{2}} \frac{d\theta}{1-\frac{1-\cos\frac{k\pi}{2^{n-1}}}{2}\sin^2 2\theta} \\
=& \frac{2}{t}\int_0^{\frac{\pi}{2}} \frac{d\theta}{1-\frac{1}{2}\left(1-\cos\frac{k\pi}{2^{n-1}}\right)\left(\frac{1-\cos 4\theta}{2}\right)} \\
=& \frac{1}{2t}\int_0^{2\pi} \frac{d\phi}{1-\frac{1-\cos\frac{k\pi}{2^{n-1}}}{4}+\frac{1}{4}\left(1-\cos\frac{k\pi}{2^{n-1}}\right)\cos\phi} \\
=& \frac{2}{t}\int_0^{2\pi} \frac{d\phi}{\left(3+\cos\frac{k\pi}{2^{n-1}}\right)+\left(1-\cos\frac{k\pi}{2^{n-1}}\right)\cos\phi} \\
=& \frac{2}{t}\frac{2\pi}{\sqrt{\left(3+\cos\frac{k\pi}{2^{n-1}}\right)^2-\left(1-\cos\frac{k\pi}{2^{n-1}}\right)^2}} \\
=& \frac{\pi\sqrt{2}}{t}\frac{1}{\sqrt{1+\cos\frac{k\pi}{2^{n-1}}}} \\
=& \frac{\pi}{t}\frac{1}{\cos\frac{k\pi}{2^n}}. \qquad (2.12)
\end{aligned}$$

From (2.7), in view of (2.12), we can conclude that

$$\int_{-\infty}^{+\infty} p_{2^n}(x,t)\,dx = 1$$

Remark 2.3. From (2.4), for $n=2$ we obtain that

$$p_4(x,t) = \frac{1}{2\pi}\left[\frac{te^{i\frac{\pi}{4}}}{x^2+(te^{i\frac{\pi}{4}})^2}+\frac{te^{-i\frac{\pi}{4}}}{x^2+(te^{-i\frac{\pi}{4}})^2}\right]$$

with Fourier transform

$$\int_{-\infty}^{+\infty} e^{i\beta x}p_4(x,t)dx = e^{-\frac{t}{\sqrt{2}}|\beta|}\cos\frac{\beta t}{\sqrt{2}}.$$

From (2.6) and (2.7) we have that

$$p_4(x,t) = \frac{t}{\sqrt{2}\pi}\frac{x^2+t^2}{x^4+t^4}. \qquad (2.13)$$

The law (2.13) has two maxima as Figure 1 shows.

Remark 2.4. For $n = 3$, from (2.6), we have that

$$p_8(x,t) = \frac{t(x^2+t^2)}{2\pi(x^8+t^8)}\left[\left(x^4+t^4+2x^2t^2\cos\frac{\pi}{4}\right)\cos\frac{3\pi}{8}+\left(x^4+t^4+2x^2t^2\cos\frac{3\pi}{4}\right)\cos\frac{\pi}{8}\right].$$

From the fact that

$$\cos\frac{3\pi}{4} = -\cos\frac{\pi}{4} \quad \text{and} \quad \cos\frac{3\pi}{8} = \sin\frac{\pi}{8}$$

we write

$$p_8(x,t) = \frac{t(x^2+t^2)}{2\pi(x^8+t^8)}\left[\left(x^4+t^4+\sqrt{2}x^2t^2\right)\sin\frac{\pi}{8}+\left(x^4+t^4-\sqrt{2}x^2t^2\right)\cos\frac{\pi}{8}\right].$$

From (2.7) we have also that

$$p_8(x,t) = \frac{t}{2\pi}\left[\frac{x^2+t^2}{x^4+t^4-\sqrt{2}x^2t^2}\sin\frac{\pi}{8}+\frac{x^2+t^2}{x^4+t^4+\sqrt{2}x^2t^2}\cos\frac{\pi}{8}\right]. \tag{2.14}$$

From (2.4) we obtain the characteristic function

$$\int_{\mathbb{R}} e^{i\beta x}p_8(x,t)dx = \frac{1}{2^2}\left[e^{-t|\beta|\cos\frac{\pi}{8}}\cos\left(t\beta\sin\frac{\pi}{8}\right)+e^{-t|\beta|\sin\frac{\pi}{8}}\cos\left(t\beta\cos\frac{\pi}{8}\right)\right].$$

The density $p_8(x,t)$ is a bimodal curve as well as $p_4(x,t)$. The maxima of $p_8(x,t)$ are heigher than those of $p_4(x,t)$ as Figure 1 shows. Also $p_{2^n}(x,t)$ displays a bimodal structure with the height peaks increasing as n increases. The form of $p_{2^n}(x,t)$ reminds the structure of densities of fractional diffusions governed by equations

$$\frac{\partial^\nu u}{\partial t^\nu} = \lambda^2\frac{\partial^2 u}{\partial x^2}$$

for $1 < \nu < 2$ (see [10]).

Remark 2.5. The result (2.7) can conveniently be rewritten as

$$p_{2^n}(x,t) = \frac{t}{\pi(x^2+t^2)}\left[\frac{1}{2^{n-1}}\sum_{\substack{k=1\\k\text{ odd}}}^{2^{n-1}-1}\frac{x^4+t^4+2x^2t^2}{x^4+t^4+2x^2t^2\cos\frac{k\pi}{2^{n-1}}}\cos\frac{k\pi}{2^n}\right]. \tag{2.15}$$

The factor in square parenthesis measures, in some sense, the disturbance of p_{2^n} on the classical Cauchy. For $n = 2$, we have in particular that

$$p_{2^2}(x,t) = \frac{t}{\pi(x^2+t^2)}\frac{1}{\sqrt{2}}\left[1+\frac{2x^2t^2}{x^4+t^4}\right] = \frac{t}{\sqrt{2}\pi}\frac{x^2+t^2}{x^4+t^4}. \tag{2.16}$$

The density (2.16) has two symmetric maxima at $x = \pm t\sqrt{\sqrt{2}-1}$ and a minimum at $x = 0$ (see Fig. 6 of Orsingher and D'Ovidio [11]). The terms

$$g_k(x,t) = \frac{x^4+t^4+2x^2t^2}{x^4+t^4+2x^2t^2\cos\frac{k\pi}{2^{n-1}}} \tag{2.17}$$

display two maxima at $x = \pm t$ with height depending on k and whose profile is depicted in Figure 2.

FIGURE 1. The profile of the functions p_4 (dotted line), formula (2.13) and p_8, formula (2.14).



FIGURE 2. The profile of the function g_k for $n = 3$ and $k = 1$ (dotted line), $k = 3$.



FIGURE 3. The profile of the functions p_{2^n}, formula (2.7), for $n = 5, 10, 15, 20$.



Remark 2.6. The density $p_{2^n}(x,t)$ can be written as

$$p_{2^n}(x,t) = \frac{t(x^2+t^2)}{2^{n-2}\pi(x^{2^n}+t^{2^n})} Q(x,t) \tag{2.18}$$

where $Q(x,t)$ is a polynomial of order $2^n - 2^2$. For $n = 2$ the function $Q(x,t)$ reduces to $\cos\frac{\pi}{4}$. For $n = 3$,

$$Q(x,t) =(x^4+t^4+\sqrt{2}x^2t^2)\sin\frac{\pi}{8} + (x^4+t^4-\sqrt{2}x^2t^2)\cos\frac{\pi}{8}.$$

The expression (2.18) shows that the probability law $p_{2^n}(x,t)$, $x \in \mathbb{R}$, $t > 0$ shares with the classical Cauchy density the property of non-existence of the mean value.

Remark 2.7. The density of the hyper Cauchy can also be presented in an alternative form by regrouping the terms in the right-hand side of (2.7) as

$$\sum_{\substack{k=1\\ \text{k odd}}}^{2^{n-2}-1} \left[\frac{\sin\frac{k\pi}{2^n}}{x^4+t^4-2x^2t^2\cos\frac{k\pi}{2^{n-1}}} + \frac{\cos\frac{k\pi}{2^n}}{x^4+t^4+2x^2t^2\cos\frac{k\pi}{2^{n-1}}} \right] \tag{2.19}$$

$$= \sum_{\substack{k=1\\ \text{k odd}}}^{2^{n-2}-1} \frac{\left(x^4+t^4+2x^2t^2\cos\frac{k\pi}{2^{n-1}}\right)\sin\frac{k\pi}{2^n} + \left(x^4+t^4-2x^2t^2\cos\frac{k\pi}{2^{n-1}}\right)\cos\frac{k\pi}{2^n}}{x^8+t^8-2x^4t^4\cos\frac{k\pi}{2^{n-2}}}.$$

For $n = 3$, from (2.19), we get again that

$$p_8(x,t) = \frac{t(x^2+t^2)}{2\pi(x^8+t^8)}\left[\left(x^4+t^4+\sqrt{2}x^2t^2\right)\sin\frac{k\pi}{8}+\left(x^4+t^4-\sqrt{2}x^2t^2\right)\cos\frac{k\pi}{8}\right].$$

Remark 2.8. The r.v.

$$W(t) = \left|C\left(t\cos\frac{\pi k}{2^n}\right) - t\sin\frac{\pi k}{2^n}\right| \tag{2.20}$$

(where $C(t)$, $t > 0$ is the Cauchy process) has probability density

$$f_k(w,t) = \frac{2t(w^2+t^2)\cos\frac{k\pi}{2^n}}{\pi(w^4+t^4+2w^2t^2\cos\frac{k\pi}{2^{n-1}})}, \quad w > 0. \tag{2.21}$$

Indeed, we have that

$$Pr\left\{W(t) < w\right\} = \int_{-w+t\sin\frac{k\pi}{2^n}}^{+w+t\sin\frac{k\pi}{2^n}} dy \frac{t\cos\frac{k\pi}{2^n}}{\pi(y^2+t^2\cos^2\frac{k\pi}{2^n})} \tag{2.22}$$

and

$$\begin{aligned}
f_k(w,t) =& \frac{d}{dw}Pr\left\{\left|C\left(t\cos\frac{\pi k}{2^n}\right) - t\sin\frac{\pi k}{2^n}\right| < w\right\}\\
=& \frac{t\cos\frac{k\pi}{2^n}}{\pi\left((w+t\sin\frac{k\pi}{2^n})^2+t^2\cos^2\frac{k\pi}{2^n}\right)} + \frac{t\cos\frac{k\pi}{2^n}}{\pi\left((-w+t\sin\frac{k\pi}{2^n})^2+t^2\cos^2\frac{k\pi}{2^n}\right)}\\
=& \frac{t\cos\frac{k\pi}{2^n}}{\pi\left(w^2+2wt\sin\frac{k\pi}{2^n}+t^2\right)} + \frac{t\cos\frac{k\pi}{2^n}}{\pi\left(w^2-2wt\sin\frac{k\pi}{2^n}+t^2\right)}\\
=& \frac{2t(w^2+t^2)\cos\frac{k\pi}{2^n}}{\pi(w^2+t^2+2wt\sin\frac{k\pi}{2^n})(w^2+t^2-2wt\sin\frac{k\pi}{2^n})}\\
=& \frac{2t(w^2+t^2)\cos\frac{k\pi}{2^n}}{\pi(w^4+t^4+2w^2t^2\cos\frac{k\pi}{2^{n-1}})}
\end{aligned}$$

because

$$2\sin^2\frac{k\pi}{2^n} = 1-\cos\frac{k\pi}{2^{n-1}}.$$

By symmetrizing (2.20) as follows

$$Z(t) = \frac{W_1(t)-W_2(t)}{2}$$

where $W_1(t), W_2(t)$ are independent copies of $W(t)$ we obtain a distribution of the form

$$h_k(w,t) = \frac{t(w^2+t^2)\cos\frac{k\pi}{2^n}}{\pi(w^4+t^4+2w^2t^2\cos\frac{k\pi}{2^{n-1}})}, \quad w \in \mathbb{R} \tag{2.23}$$

which coincides with each term of (2.15). This construction explains the reason for which each term in (2.15) has two symmetric maxima at $w = \pm t\sqrt{2\sin\frac{k\pi}{2^n}-1}$ for $k : \sin\frac{\pi k}{2^n} > \frac{1}{2}$.

FIGURE 4. The figure shows how the distribution (2.23) can be constructed from the Cauchy density by folding and symmetrizing, in the cases $n = 3$, $k = 1$ (top figures) and $k = 3$ (bottom figures). The dotted line gives the density of the folded distribution (3.13).



3. HIGHER-ORDER LAPLACE-TYPE EQUATION

Let us consider the pseudo-processes related to higher-order heat-type equations

$$\frac{\partial u}{\partial t} = c_n \frac{\partial^n u}{\partial x^n}, \quad x \in \mathbb{R},\ t > 0, \quad n > 2. \tag{3.1}$$

where $c_n = (-1)^{\frac{n}{2}+1}$ for n even and $c_n = \pm 1$ for n odd.

Pseudo-processes constructed by exploiting the sign-varying measures obtained as fundamental solutions to (3.1) have been examined in many papers since the beginning of the Sixties. A description of the procedure of construction of pseudo-processes can be found, for example in Krylov [4], Ladokhin [6], Hochberg [3], Orsingher [9], Lachal [5]. In the case where $n = 2k+1$, $c_{2k+1} = -1$, the fundamental solution to (3.1) reads

$$u_{2k+1}(x,t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-i\beta x + i(-1)^k t\beta^{2k+1}} d\beta. \tag{3.2}$$

In particular, for $k = 1$

$$u_3(x,t) = \frac{1}{\pi} \int_0^{\infty} \cos\left(\beta x + \beta^3 t\right) d\beta = \frac{1}{\sqrt[3]{3t}} Ai\left(\frac{x}{\sqrt[3]{3t}}\right) \tag{3.3}$$

where

$$Ai(x) = \frac{\sqrt{x}}{3}\left[I_{-\frac{1}{3}}\left(\frac{2}{3}x^{3/2}\right) - I_{\frac{1}{3}}\left(\frac{2}{3}x^{3/2}\right)\right]$$

is the Airy function (see for example Lebedev [7]).

In this section we study the composition of pseudo-processes with stable processes $S_\alpha(t)$, $t>0$, $\alpha \in (0,1)$ whose characteristic function reads

$$\mathbb{E}e^{i\beta S_\alpha(t)} = \exp\left(-t|\beta|^\alpha e^{-i\frac{\pi\gamma}{2}\frac{\beta}{|\beta|}}\right) = \exp\left(-\sigma\, t|\beta|^\alpha\left(1-i\theta\frac{\beta}{|\beta|}\tan\frac{\pi\alpha}{2}\right)\right) \tag{3.4}$$

where $\sigma = \cos \pi\gamma/2 > 0$ and

$$\theta = \cot\left(\frac{\pi\alpha}{2}\right)\tan\left(\frac{\pi\gamma}{2}\right).$$

The parameter γ must be chosen in such a way that $\theta \in [-1,1]$ for $\alpha \in (0,1)$. The skewness parameter $\theta = 1$ (that is $\gamma = \alpha$) corresponds to positively skewed stable distributions. For the density

$$p_\alpha(x,\gamma,t) = \frac{1}{2\pi}\int_{-\infty}^{+\infty} e^{-i\beta x}\mathbb{E}e^{i\beta S_\alpha(t)}d\beta$$

we have the scaling property

$$p_\alpha(x,\gamma,t) = \frac{1}{t^{1/\alpha}}p_\alpha\left(\frac{x}{t^{1/\alpha}},\gamma,1\right). \tag{3.5}$$

For $\alpha \in (0,1)$, we have the series representation of stable density (see [10, page 245])

$$p_\alpha(x;\gamma,1) = \frac{\alpha}{\pi}\sum_{r=0}^{\infty}(-1)^r\frac{\Gamma(\alpha(r+1))}{r!}x^{-\alpha(r+1)-1}\sin\left(\frac{\pi}{2}(\gamma+\alpha)(r+1)\right). \tag{3.6}$$

Theorem 3.1. *The composition of the pseudo-process* $X_{2k+1}(t)$, $t>0$ *with the stable process* $S_{\frac{1}{2k+1}}(t)$, $t>0$, $k \in \mathbb{N}$, *has a Cauchy probability distribution which can be written as*

$$Pr\{X_{2k+1}(S_{\frac{1}{2k+1}}(t) \in dx\}/dx = \frac{t\,\cos\frac{\pi}{2(2k+1)}}{\pi\left[\left(x+(-1)^{k+1}t\,\sin\frac{\pi}{2(2k+1)}\right)^2 + t^2\,\cos^2\frac{\pi}{2(2k+1)}\right]} \tag{3.7}$$

with $x \in \mathbb{R},\ t>0$. *The density function* (3.7) *is a solution to the higher-order Laplace equation*

$$\frac{\partial^{2k+1}u}{\partial t^{2k+1}} + \frac{\partial^{2k+1}u}{\partial x^{2k+1}} = 0, \quad x \in \mathbb{R},\ t>0 \tag{3.8}$$

Proof. For $\theta = 1$, $\alpha = \gamma = 1/2k+1$, in view of (3.4) we have that

$$\begin{aligned} U(\beta,t) &= \int_{-\infty}^{+\infty} e^{i\beta x}Pr\{X_{2k+1}(S_{\frac{1}{2k+1}}(t)) \in dx\} \\ &= \int_0^\infty Pr\{S_{\frac{1}{2k+1}}(t) \in ds\}\int_{-\infty}^{+\infty} e^{i\beta x}u_{2k+1}(x,s)\,dx \\ &= \int_0^\infty e^{is(-1)^k\beta^{2k+1}}Pr\{S_{\frac{1}{2k+1}}(t) \in ds\} \\ &= \exp\left(-t\left|(-1)^k\beta^{2k+1}\right|^{\frac{1}{2k+1}}\cos\frac{\pi}{2(2k+1)}\left(1-i\,\mathrm{sgn}\,\left((-1)^k\beta^{2k+1}\right)\tan\frac{\pi}{2(2k+1)}\right)\right) \\ &= \exp\left(-t|\beta|\left(\cos\frac{\pi}{2(2k+1)} - i(-1)^k\frac{\beta}{|\beta|}\sin\frac{\pi}{2(2k+1)}\right)\right) \end{aligned}$$

$$= \exp\left(-t|\beta|\cos\frac{\pi}{2(2k+1)} - i(-1)^k t\beta \sin\frac{\pi}{2(2k+1)}\right). \tag{3.9}$$

This is the characteristic function of a Cauchy distribution with scale parameter $t\cos\frac{\pi}{2(2k+1)}$ and location parameter $t(-1)^{k+1}\sin\frac{\pi}{2(2k+1)}$. Formula (3.9) can also be rewritten as

$$\begin{aligned} U(\beta,t) &= \exp\left(-t|\beta|\left(\cos\frac{\pi}{2(2k+1)} - i(-1)^k\frac{\beta}{|\beta|}\sin\frac{\pi}{2(2k+1)}\right)\right) \\ &= \exp\left(-t|\beta|\left(\cos\left(\frac{\pi}{2(2k+1)}(-1)^k\frac{\beta}{|\beta|}\right) - i\sin\left(\frac{\pi}{2(2k+1)}(-1)^k\frac{\beta}{|\beta|}\right)\right)\right) \\ &= \exp\left(-t|\beta|e^{-i\frac{\pi}{2(2k+1)}(-1)^k\frac{\beta}{|\beta|}}\right). \end{aligned} \tag{3.10}$$

The Fourier transform of equation (3.8) becomes

$$\frac{\partial^{2k+1}U}{\partial t^{2k+1}} + (-i\beta)^{2k+1}U = 0. \tag{3.11}$$

The derivative of order $2k+1$ of (3.10) is

$$\frac{\partial^{2k+1}U}{\partial t^{2k+1}}(\beta,t) = (-|\beta|)^{2k+1}\left(e^{-i\frac{\pi}{2(2k+1)}(-1)^k\frac{\beta}{|\beta|}}\right)^{2k+1} U(\beta,t) \tag{3.12}$$

and this shows that the Cauchy distribution (3.7) solves the higher-order Laplace equation (3.8). □

Remark 3.2. We notice that

$$\begin{aligned} \int_0^\infty Pr\{X_{2k+1}(S_{\frac{1}{2k+1}}(t)) \in dx\} =& \frac{1}{\pi}\int_{(-1)^{k+1}\tan\frac{\pi}{2(2k+1)}}^\infty \frac{dy}{1+y^2} \\ =& \frac{1}{2}\left(1+\frac{(-1)^k}{2k+1}\right) \end{aligned} \tag{3.13}$$

which is somehow in accord with Lachal [5]. The results (3.7) and (3.13) show that the mode of the Cauchy law (3.7) approaches the origin as k increases.

Let us consider the process of the form $X_3(S_{\frac{1}{3}}(t))$, $t>0$ where X_3 is a pseudo-process whose measure density is governed by the third-order heat equation

$$\frac{\partial u}{\partial t} = -\frac{\partial^3 u}{\partial x^3}, \quad x\in\mathbb{R},\ t>0 \tag{3.14}$$

and $S_{\frac{1}{3}}$ is the stable process of order 1/3. The distribution of $X_3(S_{\frac{1}{3}}(t))$, $t>0$ reads

$$Pr\{X_3(S_{\frac{1}{3}}(t))\in dx\} = dx\int_0^\infty \frac{1}{\sqrt[3]{3s}}Ai\left(\frac{x}{\sqrt[3]{3s}}\right)\frac{t}{s}\frac{1}{\sqrt[3]{3s}}Ai\left(\frac{t}{\sqrt[3]{3s}}\right)ds \tag{3.15}$$

where

$$Pr\{S_{\frac{1}{3}}(t)\in ds\} = ds\frac{t}{s}\frac{1}{\sqrt[3]{3s}}Ai\left(\frac{t}{\sqrt[3]{3s}}\right), \quad s\ge 0,\ t>0 \tag{3.16}$$

for which

$$\begin{aligned} \int_0^\infty Pr\{S_{\frac{1}{3}}(t)\in ds\} =& \int_0^\infty ds\frac{t}{s}\frac{1}{\sqrt[3]{3s}}Ai\left(\frac{t}{\sqrt[3]{3s}}\right) \\ =& (w=t/\sqrt[3]{3s}) = 3\int_0^\infty Ai(w)\,dw = 1. \end{aligned}$$

Corollary 3.3. *The law* (3.15) *solves the higher-order Laplace equation*

$$\frac{\partial^3 u}{\partial t^3} + \frac{\partial^3 u}{\partial x^3} = 0, \quad x \in \mathbb{R},\ t > 0 \tag{3.17}$$

and can be written as

$$\begin{aligned} Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\} =& \frac{dx}{\pi} \frac{\frac{\sqrt{3}}{2} t}{\left(x + \frac{t}{2}\right)^2 + \frac{3t^2}{4}} \\ =& \frac{dx}{\pi} \frac{3^{1/2}}{2} \frac{t}{x^2 + xt + t^2} \\ =& dx \frac{3^{1/2}\, t}{2\pi} \frac{x - t}{x^3 - t^3}. \end{aligned} \tag{3.18}$$

Proof. The Fourier transform of (3.15) becomes

$$\int_{-\infty}^{\infty} e^{i\beta x} Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\} = \int_0^{\infty} e^{-i\beta^3 s} Pr\{S_{\frac{1}{3}}(t) \in ds\}. \tag{3.19}$$

We show that (3.16) is a stable law of order 1/3. In view of the representation of the the Airy function (4.10) of Orsingher and Beghin [10]

$$Ai(w) = \frac{3^{-2/3}}{\pi} \sum_{k=0}^{\infty} \frac{(3^{1/3} w)^k}{k!} \sin\left(\frac{2\pi}{3}(k+1)\right) \Gamma\left(\frac{k+1}{3}\right) \tag{3.20}$$

we can write that

$$\frac{t}{s} \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{t}{\sqrt[3]{3s}}\right) = \frac{t}{3\pi s \sqrt[3]{s}} \sum_{k=0}^{\infty} \left(\frac{t}{\sqrt[3]{s}}\right)^k \frac{1}{k!} \sin\left(\frac{2\pi}{3}(k+1)\right) \Gamma\left(\frac{k+1}{3}\right)$$

We consider the series expansion (3.6) of the stable density (with $t = 1$) for which (3.4) holds true. For $\alpha = \gamma = 1/3$ (that is $\theta = +1$), $x = s/t^3$ in (3.6) we get that

$$\begin{aligned} p_{\frac{1}{3}}\left(\frac{s}{t^3}; \frac{1}{3}, 1\right) =& \frac{1}{3\pi} \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} \left(\frac{s}{t^3}\right)^{-\frac{k+1}{3} - 1} \sin\left(\frac{\pi}{3}(k+1)\right) \Gamma\left(\frac{k+1}{3}\right) \\ =& \text{(by 4.5 of [10])} \\ =& \frac{1}{3\pi} \frac{t^4}{s \sqrt[3]{s}} \sum_{k=0}^{\infty} \left(\frac{t}{\sqrt[3]{s}}\right)^k \frac{1}{k!} \sin\left(\frac{2\pi}{3}(k+1)\right) \Gamma\left(\frac{k+1}{3}\right) \\ =& t^3 \left[\frac{t}{s} \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{t}{\sqrt[3]{3s}}\right)\right] \end{aligned}$$

and thus, from (3.5), we have that

$$\frac{1}{t^3} p_{\frac{1}{3}}\left(\frac{s}{t^3}; \frac{1}{3}, 1\right) = p_{\frac{1}{3}}\left(s; \frac{1}{3}, t\right) = \frac{t}{s} \frac{1}{\sqrt[3]{3s}} Ai\left(\frac{t}{\sqrt[3]{3s}}\right), \quad s, t > 0.$$

We now evaluate the integral (3.19). We have that

$$\begin{aligned} &\int_0^{\infty} e^{-i\beta^3 s} Pr\{S_{\frac{1}{3}}(t) \in ds\} \\ =& \exp\left(-\cos\frac{\pi}{6}\, t| - \beta^3|^{\frac{1}{3}} \left(1 - i \text{ sgn } (-\beta^3) \tan\frac{\pi}{6}\right)\right) \end{aligned}$$

$$
\begin{aligned}
&= \exp\left(-\frac{\sqrt{3}}{2}t|\beta|\left(1+i\ \mathrm{sgn}\ (\beta)\frac{1}{\sqrt{3}}\right)\right)\\
&= \exp\left(-\frac{\sqrt{3}}{2}t|\beta| - i\frac{t}{2}\beta\right) \qquad (3.21)
\end{aligned}
$$

since sgn $(-\beta^3) =$ sgn $(-\beta) = -$sgn $(\beta) = -\frac{\beta}{|\beta|}$. From (3.21) we infer that

$$
\begin{aligned}
Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\} =& \frac{dx}{2\pi}\int_{-\infty}^{+\infty} e^{-i\beta x}\exp\left(-\frac{\sqrt{3}}{2}t|\beta| - i\frac{t}{2}\beta\right) d\beta \qquad (3.22)\\
=& \frac{dx}{\pi}\frac{\frac{\sqrt{3}}{2}t}{\left(x+\frac{t}{2}\right)^2+\frac{3t^2}{4}} = \frac{dx}{\pi}\frac{3^{1/2}}{2}\frac{t}{x^2+xt+t^2}\\
=& dx\frac{3^{1/2}\,t}{2\pi}\frac{x-t}{x^3-t^3}
\end{aligned}
$$

□

Remark 3.4. We observe that the r.v. $X_3(S_{\frac{1}{3}}(t))$ possesses Cauchy distribution with scale parameter $\sqrt{3}t/2$ and location parameter $-t/2$. Furthermore, it solves the third-order Laplace-type equation

$$
\frac{\partial^3 u}{\partial t^3} + \frac{\partial^3 u}{\partial x^3} = 0. \qquad (3.23)
$$

Remark 3.5. From the fact that

$$
\frac{1}{\sqrt[3]{3t}}Ai\left(\frac{x}{\sqrt[3]{3t}}\right) = \frac{1}{3\pi}\sqrt{\frac{x}{t}}K_{1/3}\left(\frac{2}{3^{3/2}}\frac{x^{3/2}}{\sqrt{t}}\right), \quad x,t>0 \qquad (3.24)
$$

we can write, for $x>0$,

$$
\begin{aligned}
Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\}/dx =& \int_0^\infty \frac{1}{3\pi}\sqrt{\frac{x}{s}}K_{1/3}\left(\frac{2}{3^{3/2}}\frac{x^{3/2}}{\sqrt{s}}\right)\frac{t}{s}\frac{1}{3\pi}\sqrt{\frac{t}{s}}K_{1/3}\left(\frac{2}{3^{3/2}}\frac{t^{3/2}}{\sqrt{s}}\right) ds \qquad (3.25)\\
=& \frac{2\sqrt{xt^3}}{3^2\pi^2}\int_0^\infty s\,K_{1/3}\left(\frac{2x^{3/2}}{3^{3/2}}s\right)K_{1/3}\left(\frac{2t^{3/2}}{3^{3/2}}s\right) ds. \qquad (3.26)
\end{aligned}
$$

In view of (see [2, formula 6.521])

$$
\int_0^\infty s\,K_\nu(ys)\,K_\nu(zs)\,ds = \frac{\pi(yz)^{-\nu}(y^{2\nu}-z^{2\nu})}{2\sin\pi\nu\,(y^2-z^2)}, \quad \Re\{y+z\}>0,\ |\Re\{\nu\}|<1
$$

we get that

$$
Pr\{X_3(S_{\frac{1}{3}}(t)) \in dx\} = dx\,\frac{3^{1/2}\,t}{2\pi}\frac{x-t}{x^3-t^3}, \quad x,t>0 \qquad (3.27)
$$

which coincides with (3.18).

The Cauchy densities pertaining to the composition $X_{\frac{1}{2k+1}}(S_{\frac{1}{2k+1}}(t))$, $t>0$, solve also a second-order p.d.e. as we show in the next theorem.

FIGURE 5. The profile of the function (3.18).



Theorem 3.6. *The Cauchy densities*

$$f(x,t;m) = \frac{1}{\pi} \frac{t \cos \frac{\pi}{2m}}{(x + t \sin \frac{\pi}{2m})^2 + t^2 \cos^2 \frac{\pi}{2m}}, \quad m \in \mathbb{N} \tag{3.28}$$

satisfy the following second-order equation

$$\frac{\partial^2 f}{\partial t^2} + \frac{\partial^2 f}{\partial x^2} = 2 \sin \frac{\pi}{2m} \frac{\partial^2 f}{\partial x \partial t}, \quad x \in \mathbb{R},\ t > 0. \tag{3.29}$$

Proof. It is convenient to write (3.28) as a composed function

$$f(u,v) = \frac{1}{\pi} \frac{u}{u^2 + v^2}$$

where

$$u = t \cos \frac{\pi}{2m}, \qquad v = x + t \sin \frac{\pi}{2m}.$$

Since

$$\begin{aligned}
\frac{\partial f}{\partial t} &= \cos \frac{\pi}{2m} \frac{\partial f}{\partial u} + \sin \frac{\pi}{2m} \frac{\partial f}{\partial v} \\
\frac{\partial^2 f}{\partial t^2} &= \cos^2 \frac{\pi}{2m} \frac{\partial^2 f}{\partial u^2} + 2 \cos \frac{\pi}{2m} \sin \frac{\pi}{2m} \frac{\partial^2 f}{\partial u \partial v} + \sin^2 \frac{\pi}{2m} \frac{\partial^2 f}{\partial v^2} \\
\frac{\partial f}{\partial x} &= \frac{\partial f}{\partial v} \quad \text{and} \quad \frac{\partial^2 f}{\partial x^2} = \frac{\partial^2 f}{\partial v^2}
\end{aligned}$$

and

$$\frac{\partial^2 f}{\partial u^2} + \frac{\partial^2 f}{\partial v^2} = 0$$

we have that

$$\begin{aligned}
\frac{\partial^2 f}{\partial t^2} + \frac{\partial^2 f}{\partial x^2} =& \cos^2 \frac{\pi}{2m} \frac{\partial^2 f}{\partial u^2} + \frac{\partial^2 f}{\partial v^2} + 2 \sin \frac{\pi}{2m} \cos \frac{\pi}{2m} \frac{\partial^2 f}{\partial u \partial v} + \sin^2 \frac{\pi}{2m} \frac{\partial^2 f}{\partial v^2} \\
=& \frac{\partial^2 f}{\partial v^2} \left[1 - \cos^2 \frac{\pi}{2m} + \sin^2 \frac{\pi}{2m} \right] + 2 \sin \frac{\pi}{2m} \cos \frac{\pi}{2m} \frac{\partial^2 f}{\partial u \partial v} \\
=& 2 \sin \frac{\pi}{2m} \frac{\partial}{\partial v} \left[\sin \frac{\pi}{2m} \frac{\partial f}{\partial v} + \cos \frac{\pi}{2m} \frac{\partial f}{\partial u} \right] \\
=& 2 \sin \frac{\pi}{2m} \frac{\partial}{\partial x} \frac{\partial f}{\partial t}
\end{aligned}$$

□

Remark 3.7. The characteristic function of (3.28) is

$$\int_{-\infty}^{+\infty} e^{i\beta x} f(x,t;m)dx = e^{-t|\beta|\cos\frac{\pi}{2m} - i\beta t \sin\frac{\pi}{2m}}$$

and can be obtained by considering the bounded solution to the Fourier transform of (3.29)

$$\frac{d^2F}{dt^2} + 2i\beta \sin\frac{\pi}{2m}\frac{dF}{dt} - \beta^2 F = 0.$$

For the even-order Laplace equations we have the following result.

Theorem 3.8. *The solution to the higher-order Laplace-type equation*

$$\frac{\partial^{2n}u}{\partial t^{2n}} = -\frac{\partial^{2n}u}{\partial x^{2n}}, \quad x\in\mathbb{R},\ t>0 \tag{3.30}$$

subject to the initial conditions

$$\begin{cases} u(x,0) = \delta(x) \\ \frac{\partial^k u}{\partial t^k}(x,t)\Big|_{t=0^+} = \frac{(-1)^k\, k!}{\pi|x|^{k+1}}\cos\frac{\pi(k+1)}{2}, \quad 0<k<2n \end{cases} \tag{3.31}$$

is the classical Cauchy distribution given by

$$u(x,t) = Pr\{X_{2n}(S_{\frac{1}{2n}}(t)) \in dx\}/dx = \frac{t}{\pi(x^2+t^2)}, \quad x\in\mathbb{R},\ t>0 \tag{3.32}$$

where $X_{2n}(t)$, $t>0$ is a pseudo-process such that

$$\mathbb{E}e^{i\beta X_{2n}(t)} = e^{-t\beta^{2n}}.$$

Proof. The pseudo-process $X_{2n}(t)$, $t>0$ related to the equation

$$\frac{\partial u}{\partial t} = (-1)^{n+1}\frac{\partial^{2n}u}{\partial t^{2n}}$$

has fundamental solution whose Fourier transform reads

$$\int_{-\infty}^{+\infty} e^{i\beta x}u(x,t)dx = e^{-t\beta^{2n}}.$$

If $S_{\frac{1}{2n}}(t)$, $t>0$ is a stable subordinator with Laplace transform

$$\mathbb{E}\exp\left(-\lambda S_{\frac{1}{2n}}(t)\right) = \exp\left(-t\lambda^{\frac{1}{2n}}\right), \quad \lambda>0,\ t>0 \tag{3.33}$$

the characteristic function of $X_{2n}(S_{\frac{1}{2n}}(t))$, $t>0$ becomes

$$\begin{aligned}\int_{-\infty}^{+\infty} e^{i\beta x}Pr\{X_{2n}(S_{\frac{1}{2n}}(t))\in dx\} &= \int_0^\infty e^{-s\beta^{2n}}Pr\{S_{\frac{1}{2n}}(t)\in ds\} \\ &= \exp\left(-t|\beta|e^{i\frac{\pi r}{n}}\right), \quad r=0,1,\dots,2n-1 \end{aligned}\tag{3.34}$$

For $r=0$, we have the characteristic function of the Cauchy symmetric law. For $r\neq 0$ and $n\le r\le 2n-1$ we have a function which is not absolutely integrable and, for $0<r<n-1$ is not a characteristic function (but can be regarded as a Cauchy r.v. at a complex time). The functions

$$F_r(\beta,t) = e^{-t|\beta|e^{i\frac{\pi r}{2n}}}$$

for all $0 \leq r \leq 2n-1$ are solutions to

$$\frac{\partial^{2n} F_r}{\partial t^{2n}} = (-1)^{n+1} F_r.$$

We now check that for $0 \leq k \leq 2n-1$ the initial conditions (3.31) are verified by the Cauchy distribution. Indeed,

$$\begin{aligned}\frac{\partial^k u}{\partial t^k}(x,t)\Big|_{t=0} =& \frac{\partial^k}{\partial t^k}\left(\frac{1}{2\pi}\int_{-\infty}^{+\infty} e^{-i\beta x} e^{-t|\beta|} d\beta\right)\Bigg|_{t=0} \\ =& \frac{1}{2\pi}\int_{-\infty}^{+\infty} e^{-i\beta x}(-1)^k|\beta|^k d\beta \\ =& \frac{(-1)^k k!}{\pi|x|^{k+1}}\cos\left(\frac{\pi(k+1)}{2}\right).\end{aligned}$$

□

Remark 3.9. We notice that for $n=1$ the problem above becomes

$$\frac{\partial^2 u}{\partial t^2} = -\frac{\partial^2 u}{\partial x^2}, \quad x \in \mathbb{R},\ t > 0$$

subject to the initial conditions

$$\begin{cases} u(x,0) = \delta(x) \\ \frac{\partial u}{\partial t}(x,t)\Big|_{t=0^+} = \frac{-1}{\pi|x|^2}\cos\pi \end{cases}$$

which is in accord with

$$\frac{\partial}{\partial t}\frac{t}{\pi(x^2+t^2)}\Big|_{t=0^+} = \frac{1}{\pi x^2}.$$

The connection between wave equations and the composition of two independent Cauchy processes $C^1(|C^2(t)|)$, $t>0$ has been investigated in D'Ovidio and Orsingher [1] and more general results involving the Cauchy process have been presented in Nane [8].

Remark 3.10. We finally notice that the equation

$$\frac{\partial^6 u}{\partial t^6} + \frac{\partial^6 u}{\partial x^6} = 0 \tag{3.35}$$

can be decoupled as

$$\left(\frac{\partial^3}{\partial t^3} + i\frac{\partial^3}{\partial x^3}\right)\left(\frac{\partial^3}{\partial t^3} - i\frac{\partial^3}{\partial x^3}\right) u = 0. \tag{3.36}$$

Form the Corollary 3.3, the solution to (3.36) can be therefore written as

$$\begin{aligned} u(x,t) =& \frac{1}{2\pi}\left[\frac{\frac{\sqrt{3}}{2} t e^{i\frac{\pi}{6}}}{\left(x + \frac{te^{i\frac{\pi}{6}}}{2}\right)^2 + \frac{3}{4}t^2 e^{i\frac{\pi}{3}}} + \frac{\frac{\sqrt{3}}{2} t e^{-i\frac{\pi}{6}}}{\left(x + \frac{te^{-i\frac{\pi}{6}}}{2}\right)^2 + \frac{3}{4}t^2 e^{-i\frac{\pi}{3}}}\right] \\ =& \frac{\sqrt{3}}{2^2\pi} t \left[\frac{e^{i\frac{\pi}{6}}\left(x^2 + \frac{t}{4}e^{-i\frac{\pi}{3}} + xte^{-i\frac{\pi}{6}} + \frac{3}{4}t^2 e^{-i\frac{\pi}{3}}\right)}{\left(x^2 + \frac{t}{4}e^{-i\frac{\pi}{3}} + xte^{-i\frac{\pi}{6}} + \frac{3}{4}t^2 e^{-i\frac{\pi}{3}}\right)\left(x^2 + \frac{t}{4}e^{i\frac{\pi}{3}} + xte^{i\frac{\pi}{6}} + \frac{3}{4}t^2 e^{i\frac{\pi}{3}}\right)}\right. \\ &\left. + \frac{e^{-i\frac{\pi}{6}}\left(x^2 + \frac{t}{4}e^{i\frac{\pi}{3}} + xte^{i\frac{\pi}{6}} + \frac{3}{4}t^2 e^{i\frac{\pi}{3}}\right)}{\left(x^2 + \frac{t}{4}e^{-i\frac{\pi}{3}} + xte^{-i\frac{\pi}{6}} + \frac{3}{4}t^2 e^{-i\frac{\pi}{3}}\right)\left(x^2 + \frac{t}{4}e^{i\frac{\pi}{3}} + xte^{i\frac{\pi}{6}} + \frac{3}{4}t^2 e^{i\frac{\pi}{3}}\right)}\right] \end{aligned}$$

$$
\begin{aligned}
&=\frac{\sqrt{3}}{2^2\pi} t \frac{(x^2+t^2)\cos\frac{\pi}{6}+xt}{\left(x^2+te^{-i\frac{\pi}{3}}+xte^{-i\frac{\pi}{6}}+\right)\left(x^2+te^{i\frac{\pi}{3}}+xte^{i\frac{\pi}{6}}\right)} \\
&=\frac{\sqrt{3}}{2^2\pi} t \frac{(x^2+t^2)\cos\frac{\pi}{6}+xt}{\left(x^2+t^2+xt\cos\frac{\pi}{6}\right)^2+2x^2t^2\cos\frac{\pi}{3}}. \qquad (3.37)
\end{aligned}
$$

Equation (3.36) is satisfied by the Cauchy density and therefore by the probability law (3.37) which however is no longer a Cauchy distribution but is unimodal and asymmetric.

References

[1] M. D'Ovidio and E. Orsingher. Composition of processes and related partial differential equations. *J. Theor. Probab.*, 24:342 – 375, 2011.

[2] I. S. Gradshteyn and I. M. Ryzhik. *Table of integrals, series and products.* Academic Press, 2007. Seventh edition.

[3] K. J. Hochberg. A signed measure on path space related to Wiener measure. *Ann. Probab.*, 6:433 – 458, 1978.

[4] V. Yu. Krylov. Some properties of the distribution corresponding to the equation $\frac{\partial u}{\partial t} = (-1)^{q+1}\frac{\partial u^{2q}}{\partial x^{2q}}$. *Soviet Math. Dokl.*, 1:760 – 763, 1960.

[5] A. Lachal. Distributions of sojourn time, maximum and minimum for pseudo-processes governed by higher-order heat-type equations. *Elect. J. Probab.*, 8 (20):1 – 53, 2003.

[6] V.I. Ladokhin. On the measure on functional spaces corresponding to complex and diffusion coefficients. *Uchenye Zapiski Kazan Univ.*, 123:36 – 42, 1963. in Russian.

[7] N. N. Lebedev. *Special functions and their applications.* Dover, New York, 1972.

[8] E. Nane. Higher order pde's and iterated processes. *Trans. Amer. Math. Soc.*, 360(5):2681 – 2692, 2008.

[9] E. Orsingher. Processes governed by signed measures connected with third-order "heat-type" equations. *Litovsk. Mat. Sb.*, 31(2):323 – 336, 1991. Translation in Lithuanian Math. J. 31 (1991), no. 2, 220 - 231 (1992).

[10] E. Orsingher and L. Beghin. Fractional diffusion equations and processes with randomly varying time. *Ann. Probab.*, 37:206 – 249, 2009.

[11] E. Orsingher and M. D'Ovidio. Vibrations and fractional vibrations of rods, plates and Fresnel pseudo-processes. *J. Stat. Phys.*, 145:143 – 174, 2011. Published online: 01 September 2011.

DEPARTMENT OF STATISTICAL SCIENCES, SAPIENZA UNIVERSITY OF ROME
E-mail address: `enzo.orsingher@uniroma1.it`

DEPARTMENT OF STATISTICAL SCIENCES, SAPIENZA UNIVERSITY OF ROME
E-mail address: `mirko.dovidio@uniroma1.it`